

FORM PTO-1390
(REV 10-95)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. §371**

SCH 1799

U.S. APPLICATION NO. (If known, see 37 CFR §1.5)

09/787394

INTERNATIONAL APPLICATION NO

INTERNATIONAL FILING DATE

PCT/EP99/07088

16 SEPTEMBER 1999

PRIORITY DATE CLAIMED

18 SEPTEMBER 1998

TITLE OF INVENTION

NEAR INFRARED FLUORESCENT CONTRAST AGENT AND FLUORESCENCE IMAGING

APPLICANT(S) FOR DO/EO/US

MIWA, Naoto, et al.





Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. §371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. §371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. §371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. §371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. §371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. §371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. §371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. §371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. §371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. §371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 C.F.R. §§1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 C.F.R. §§3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
 - ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☐ Other items or information:

U.S. APPLICATION NO. (if known, see 37 CFR §1.5) 097787394		INTERNATIONAL APPLICATION NO. PCT/EP99/07088		ATTORNEY'S DOCKET NUMBER SCH 1799	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR §1.492 (a) (1) - (5)): Search Report has been prepared by the EPO or JPO..... \$860.00 International preliminary examination fee paid to USPTO (37 CFR §1.482)..... \$690.00 No international preliminary examination fee paid to USPTO (37 CFR §1.482) but international search fee paid to USPTO (37 CFR §1.445(a)(2))..... \$710.00 Neither international preliminary examination fee (37 CFR §1.482) nor international search fee (37 CFR §1.445(a)(2)) paid to USPTO..... \$1000.00 International preliminary examination fee paid to USPTO (37 CFR §1.482) and all claims satisfied provisions of PCT Article 33(2)-(4)..... \$100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 C.F.R. §1.492(e)). <input type="checkbox"/> 20 <input type="checkbox"/> 30					
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	28 - 20 =	8	x \$ 18.00	\$144.00	
Independent claims	3 - 3 =	0	x \$ 80.00	\$0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$ 270.00		
TOTAL OF ABOVE CALCULATIONS =				\$1,004.00	
Reduction of 1/2 for filing by small entity, if applicable. A Verified Small Entity Statement must also be filed (Note 37 C.F.R. §§1.9, 1.27, 1.28).					
SUBTOTAL =				\$1,004.00	
Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 C.F.R. §1.492(f)). <input type="checkbox"/> 20 <input type="checkbox"/> 30					
TOTAL NATIONAL FEE =				\$1,004.00	
Fee for recording the enclosed assignment (37 C.F.R. §1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 C.F.R. §§3.28, 3.31). \$40.00 per property.					
TOTAL FEES ENCLOSED =				\$1,004.00	
				Amount to be refunded:	
				charged:	
a. <input checked="" type="checkbox"/> A check in the amount of <u>\$1,004.00</u> to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. <u>13-3402</u> in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>13-3402</u> . A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 C.F.R. §§1.494 or 1.495 has not been met, a petition to revive (37 C.F.R. §1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: Customer Number 23,599 <div style="text-align: center;">  23599 PATENT TRADEMARK OFFICE </div>					
Filed: 19 MARCH 2001 AJZ:jmm			SIGNATURE  NAME <u>Anthony J. Zelano</u>		
			27,969 REGISTRATION NUMBER		

IN THE UNITED STATES DESIGNATED/ELECTED OFFICE

International Application No. : PCT/EP99/07088
International Filing Date : 16 SEPTEMBER 1999
Priority Date(s) Claimed : 18 SEPTEMBER 1998
Applicant(s) (DO/EO/US) : MIWA, Naoto, et al.
Title: NEAR INFRARED FLUORESCENT CONTRAST AGENT AND
FLUORESCENCE IMAGING

PRELIMINARY AMENDMENT

Commissioner for Patents
Washington, D.C. 20231

SIR:

Prior to calculating the national fee, and prior to examination in the National Phase of the above-identified International application, please amend as follows:

IN THE CLAIMS:

3. (Amended) The near infrared fluorescent contrast agent of claim 1, wherein, in the formula [I], r is 1.
4. (Amended) The near infrared fluorescent contrast agent of claim 1, wherein 4 or more sulfonic acid groups are contained in a molecule.
5. (Amended) The near infrared fluorescent contrast agent of claim 1, wherein 10 or less sulfonic acid groups are contained in a molecule.
6. (Amended) The near infrared fluorescent contrast agent of claim 1, wherein 8 or less sulfonic acid groups are contained in a molecule.
7. (Amended) The near infrared fluorescent contrast agent of claim 1, wherein the pharmaceutically acceptable salt is a sodium salt.

8. (Amended) The near infrared fluorescent contrast agent of claim 1, which is for tumor imaging.

9. (Amended) The near infrared fluorescent contrast agent of claim 1, which is for angiography.

18. (Amended) The sodium salt of claim 10, comprising 4 or more sulfonic acid groups in a molecule.

19. (Amended) The sodium salt of claim 10, comprising 10 or less sulfonic acid groups in a molecule.


20. (Amended) The sodium salt of claim 10, comprising 8 or less sulfonic acid groups in a molecule.

21. (Amended) A near infrared fluorescent contrast agent comprising the sodium salt of claim 10.

REMARKS

The purpose of this Preliminary Amendment is to eliminate multiple dependent claims in order to avoid the additional fee. Applicants reserve the right to reintroduce claims to canceled combined subject matter.

Respectfully submitted,



Anthony J. Zelano, Reg. No. 27,969
Attorney for Applicants
MILLEN, WHITE, ZELANO & BRANIGAN, P.C.
Arlington Courthouse Plaza 1
2200 Clarendon Boulevard, Suite 1400
Arlington, VA 22201
Direct Dial: 703-812-5311
Facsimile: 703-243-6410
Email: zelano@mwzb.com

AJZ:jmm

09787394-051601

VERSION WITH MARKINGS TO SHOW CHANGES MADE

Claims 3-9 and 18-21 have been amended as follows:

3. (Amended) The near infrared fluorescent contrast agent of claim 1 ~~or claim 2~~, wherein, in the formula [I], r is 1.

4. (Amended) The near infrared fluorescent contrast agent of ~~any of claims 1 to 3~~, claim 1, wherein 4 or more sulfonic acid groups are contained in a molecule.

5. (Amended) The near infrared fluorescent contrast agent of ~~any of claims 1 to 4~~, claim 1, wherein 10 or less sulfonic acid groups are contained in a molecule.

6. (Amended) The near infrared fluorescent contrast agent of ~~any of claims 1 to 4~~, claim 1, wherein 8 or less sulfonic acid groups are contained in a molecule.

7. (Amended) The near infrared fluorescent contrast agent of ~~any of claims 1 to 6~~, claim 1, wherein the pharmaceutically acceptable salt is a sodium salt.

8. (Amended) The near infrared fluorescent contrast agent of ~~any of claims 1 to 7~~, claim 1, which is for tumor imaging.

9. (Amended) The near infrared fluorescent contrast agent of ~~any of claims 1 to 7~~, claim 1, which is for angiography.

18. (Amended) The sodium salt of ~~any of claims 10, 11, 13, 14 and 15~~, claim 10, comprising 4 or more sulfonic acid groups in a molecule.

19. (Amended) The sodium salt of ~~any of claims 10, 11, 13, 14, 15 and~~

09/787394-051001

~~18,claim 10,~~ comprising 10 or less sulfonic acid groups in a molecule.

20. (Amended) The sodium salt of any of ~~claims 10, 11, 13, 14, 15 and~~
~~18,claim 10,~~ comprising 8 or less sulfonic acid groups in a molecule.

21. (Amended) A near infrared fluorescent contrast agent comprising the
sodium salt of any of ~~claims 10 to 20.~~claim 10.

09750"4628260

NEAR INFRARED FLUORESCENT CONTRAST AGENT ANDFLUORESCENCE IMAGING

TECHNICAL FIELD OF THE INVENTION

The present invention relates to a near infrared fluorescent contrast agent and fluorescence imaging using said contrast agent.

In treating diseases, it is vital to detect morphological and functional changes caused by the disease in the living body at an early stage of the disease. Especially when treating cancer, the site and size of tumor are significant determinant factors of the effective treatment design. Known methods for this purpose include biopsy by puncture and the like, and imaging diagnoses such as X-ray imaging, MRI, ultrasound imaging and the like. Biopsy is effective for definitive diagnosis, but at the same time it places great burden on test subjects and is not suitable for tracking time-course changes in lesions. X-ray imaging and MRI inevitably expose test subjects to radiation and magnetic waves. In addition, conventional imaging diagnoses as mentioned above require complicated operation and a long time for measurement and diagnosis. A large apparatus used for this purpose also makes it difficult to apply these methods during operation.

One of the image diagnoses is fluorescence imaging (Lipsch R. L. et al., *J. Natl. Cancer Inst.*, 26, 1-11 (1961)). This method uses, as a contrast agent, a substance that emits fluorescence upon exposure to an excitation light having a specific wavelength. Thus, a body is exposed to an excitation light from outside the body and the fluorescence emitted from the fluorescent contrast agent in the body is detected.

Such fluorescent contrast agent may be, for example, a porphyrin compound that accumulates in tumor and is used for photodynamic therapy (PDT), such as haematoporphyrin. Other examples include photophrin and benzoporphyrin

09787394-054001

(see Lipsbn R. L. et al., *supra*, Meng T. S. et al., *SPIE*, 1641, 90-98 (1992), WO 84/04665 and the like). These compounds are originally used for PDT and have phototoxicity, since this is what PDT requires. Consequently, these are not desirable diagnostic agents.

Meanwhile, retinal circulatory microangiography using a known fluorescent dye, such as fluorescein, fluorescamin and riboflavin, has been known (U.S. Patent No. 4945239). These fluorescent dyes emit fluorescence in a visible light region of 400-600 nm. In this region, the light transmission through living tissue is very low, so that the detection of lesions in the deep part of a body is nearly impossible.

In addition, the use, as fluorescent contrast agent, of cyanine compounds inclusive of indocyanine green (hereinafter to be abbreviated as ICG), which are used to determine liver function and cardiac output, has been documented (Haglund M. M. et al., *Neurosurgery*, 35, 930 (1994), Li, X. et al., *SPIE*, 2389, 789-797 (1995)). Cyanine compounds show absorbance in a near infrared light region (700-1300 nm).

Near infrared light shows high transmission through living tissues and can pass through a skull of about 10 cm in size. Because of this, it has been increasingly attracting attention in clinical medicine. For example, optical CT technique using optical transmission of medium has been drawing attention in the clinical field as a new technology. This is because near infrared light can pass through living body and can be used for monitoring oxygen concentration and circulation in the living body.

Cyanine compounds emit fluorescence in the near infrared region. The fluorescence in this region can pass through living tissues and offers the potential for a fluorescent contrast agent. Various cyanine compounds have been developed in recent years and tried as fluorescent contrast agents (WO96/17628, WP97/13490 and the like). However, an agent having sufficient

09707394 "051604
T09T50 45E 28760

solubility in water and safety to living body, as well as capability of distinguishing normal tissues from diseased tissues (selectivity for imaging target site), is not in existence.

SUMMARY OF THE INVENTION


It is therefore an object of the present invention to provide a fluorescent contrast agent. The inventive agent is low toxic and has a superior solubility in water. In addition, it emits fluorescence in a near infrared region that can pass through living tissues, and permits specific imaging of tumor and/or blood vessel.

Another object of the present invention is to provide a method of fluorescence imaging using said near infrared fluorescent contrast agent.

The present invention is predicated on the finding that introduction of three or more sulfonic acid groups into a cyanine dye compound results in the provision of a fluorescent contrast agent having a high solubility in water. It has been also found that a method of fluorescence imaging can be established when this contrast agent is used.


Thus, the present invention provides the following.

(1) A near infrared fluorescent contrast agent comprising a compound having three or more sulfonic acid groups in a molecule, which is represented by the formula [I]




wherein R^1 and R^2 are the same or different and each is a substituted or unsubstituted alkyl; Z^1 and Z^2 are each nonmetallic atoms necessary for forming a substituted or unsubstituted condensed benzo ring or condensed

naphtho ring; r is 0, 1 or 2; $L^1 - L^7$ are the same or different and each is a substituted or unsubstituted methine, provided that when r is 2, L^6 and L^7 that occur in duplicate are the same or different; and
 X and Y are the same or different and each is a group of the formula








wherein R^3 and R^4 are the same or different and each is substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.

- (2) The near infrared fluorescent contrast agent of (1) above, which is free of a carboxylic acid group in a molecule.
- (3) The near infrared fluorescent contrast agent of (1) or (2) above, wherein, in the formula [I], r is 1.
- (4) The near infrared fluorescent contrast agent of any of (1) to (3) above, wherein 4 or more sulfonic acid groups are contained in a molecule.
- (5) The near infrared fluorescent contrast agent of any of (1) to (4) above, wherein 10 or less sulfonic acid groups are contained in a molecule.
- (6) The near infrared fluorescent contrast agent of any of (1) to (4) above, wherein 8 or less sulfonic acid groups are contained in a molecule.
- (7) The near infrared fluorescent contrast agent of any of (1) to (6) above, wherein the pharmaceutically acceptable salt is a sodium salt.
- (8) The near infrared fluorescent contrast agent of any of (1) to (7) above, that is for tumor imaging and/or angiography.
- (9) A sodium salt of a compound of the formula [II] having three or more sulfonic acid groups in a molecule




wherein R^1 , R^2 , $L^1 - L^7$, X and Y are as defined above, and R^5 to R^{16} are the same or different and each is a hydrogen atom, a sulfonic acid group, a carboxyl group, a hydroxyl group, an alkyl(sulfoalkyl)amino group, a bis(sulfoalkyl)amino group, a sulfoalkoxy group, a (sulfoalkyl)sulfonyl group or a (sulfoalkyl)aminosulfonyl group, exclusive of the groups of the following formulas






(10) The sodium salt of (9) above, wherein, in the formula [II], R^1 and R^2 are each a lower alkyl having 1 to 5 carbon atoms substituted by a sulfonic acid




group and X and Y are the same or different and each is a group of the formula wherein R^{17} and R^{18} are unsubstituted lower alkyl having 1 to 5 carbon atoms.

(11) The sodium salt of (10) above, having the formula




(12) A sodium salt of a compound of the formula [III-1] having three or more








sulfonic acid groups in a molecule

wherein L^1-L^7 are as defined above, R^{19} and R^{20} are lower alkyl having 1 to 5 carbon atoms and substituted by sulfonic acid group, $R^{21} - R^{28}$ are the same or different and each is hydrogen atom, sulfonic acid group, carboxyl group, hydroxyl group, alkyl(sulfoalkyl)amino group, bis(sulfoalkyl)amino group,




sulfoalkoxy group, (sulfoalkyl)sulfonyl group or (sulfoalkyl)aminosulfonyl group, and X' and Y' are the same or different and each is a group of the formula wherein R¹⁷ and R¹⁸ are as defined above, exclusive of the groups of the following formulas






and



(13) The sodium salt of (12) above, wherein, in the formula [III-1], L_4 is methine substituted by alkyl having 1 to 4 carbon atoms.


(14) The sodium salt of (12) above, which is a sodium salt of the compound of the formula [III-2] having three or more sulfonic acid groups in a molecule




wherein $\text{R}^{19} - \text{R}^{28}$, X' and Y' are as defined above, Z^3 is a non-metallic atom group necessary for forming a 5- or 6-membered ring and A is hydrogen atom

or a monovalent group.

(15) The sodium salt of (14) above, having the formula



(16) The sodium salt of (12) above, having the formula



(17) The sodium salt of any of (9), (10), (12), (13) and (14) above, comprising 4 or more sulfonic acid groups in a molecule.

(18) The sodium salt of any of (9), (10), (12), (13), (14) and (17) above, comprising 10 or less sulfonic acid groups in a molecule.


(19) The sodium salt of any of (9), (10), (12), (13), (14) and (17) above, comprising 8 or less sulfonic acid groups in a molecule.


(20) A near infrared fluorescent contrast agent comprising the sodium salt of any of (9) to (19) above.

(21) The near infrared fluorescent contrast agent of (20) above, which is for tumor imaging and/or angiography.


(22) A method of fluorescence imaging comprising introducing the near infrared fluorescent contrast agent of (1) above into a living body, exposing the body to an excitation light, and detecting near infrared fluorescence from the contrast agent.

(23) The sodium salt of (9) above, which is at least one member selected from the group consisting of the compounds of the following formulas













and











(24) The sodium salt of (12) above, which is at least one member selected from the group consisting of the compounds of the following formulas










14
































(25) The near infrared fluorescent contrast agent of (1) above, comprising at





least one compound selected from the group consisting of the compounds of the following formulas

























































and



(26) The sodium salt of (14) above, wherein the monovalent group of A is substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted aralkyl, lower alkoxy, optionally substituted substituted amino, alkylcarbonyloxy, substituted or unsubstituted alkylthio, substituted or unsubstituted arylthio, cyano, nitro or halogen atom.

10 **BRIEF DESCRIPTION OF THE DRAWINGS**

Figs. 1 to 4 are photographs showing fluorescence imaging at 24 hours after administration of the compound, wherein administered were A:ICG (5 mg/kg), B:NK-1967 (5 mg/kg), C:compound (29) (0.5 mg/kg) and D:compound (6) K salt (5 mg/kg).

Fig. 5 is a photograph showing fluorescence imaging at 24 hours after administration of the compound, wherein administered was E:compound (31) (5 mg/kg).

Figs. 6 to 9 are photographs showing fluorescence imaging at 20 seconds and 5 minutes after administration of the compound (5 mg/kg), wherein administered were A:ICG (20 seconds later), B:ICG (5 minutes later), C:compound (29) (20 seconds later) and D:compound (29) (5 minutes later).

Fig. 10 is a graph showing the concentration of the compound in plasma at 0.5, 1, 4 and 24 hours after administration of the compound, wherein the axis of ordinates is concentration ($\mu\text{g/ml}$) of the compound in plasma at each time point.

Fig. 11 is a chart showing the infrared absorption spectrum of compound (29).

Fig. 12 is a chart showing the infrared absorption spectrum of compound (31).

5

Fig. 13 is a chart showing the infrared absorption spectrum of compound (6).

Fig. 14 is a chart showing the infrared absorption spectrum of compound (54).

10 DETAILED DESCRIPTION OF THE INVENTION


The terms used in the present specification are defined in the following.

15 The near infrared fluorescent contrast agent in the present invention means a contrast agent that emits fluorescence in a near infrared region.

In the present invention, the sulfonic acid group may mean sulfonate ($-\text{SO}_3^-$) when said sulfonic acid group is used to form an inner salt.

In the present invention, preferable X and Y are of the following formula

20



25 wherein R^3 and R^4 are the same or different and each is substituted or unsubstituted alkyl.

The alkyl of "substituted or unsubstituted alkyl" at R^1 , R^2 , R^3 and R^4 is preferably linear or branched lower alkyl having 1 to 5 carbon atoms, such as methyl, ethyl, 30 propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, tert-pentyl, 2-methylpropyl, 1,1-dimethylpropyl and the like. The substituent may be, for example, sulfonic acid group, carboxyl, hydroxy and the like. Examples of substituted alkyl include hydroxymethyl, 1-hydroxyethyl, 2-


hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 4-hydroxybutyl, carboxymethyl, carboxyethyl, carboxybutyl, sulfomethyl, 2-sulfoethyl, 3-sulfopropyl, 4-sulfobutyl and the like. Preferred R^1 and R^2 are lower alkyl having 1 to 5 carbon atoms that is substituted by sulfonic acid group (e.g., 2-sulfoethyl, 3-sulfopropyl, 4-sulfobutyl and the like), and R^3 and R^4 are unsubstituted lower alkyl having 1 to 5 carbon atoms (e.g., methyl, ethyl and the like).

The unsubstituted lower alkyl having 1 to 5 carbon atoms at R^{17} and R^{18} is exemplified by those mentioned above with regard to the alkyl of "substituted or unsubstituted alkyl" at R^1 , R^2 , R^3 and R^4 .

The alkyl group of the lower alkyl having 1 to 5 carbon atoms that is substituted by sulfonic acid group at R^{19} and R^{20} is exemplified by those mentioned above with regard to the alkyl of "substituted or unsubstituted alkyl" at R^1 , R^2 , R^3 and R^4 , and examples of the substituted lower alkyl having 1 to 5 carbon atoms include 2-sulfoethyl, 3-sulfopropyl and 4-sulfobutyl.

The alkyl moiety of alkyl(sulfoalkyl)amino group, bis(sulfoalkyl)amino group, sulfoalkoxy group, (sulfoalkyl)sulfonyl group and (sulfoalkyl)aminosulfonyl group at R^{21} - R^{28} is preferably a linear or branched lower alkyl having 1 to 5 carbon atoms, which is exemplified by those mentioned above with regard to the alkyl of "substituted or unsubstituted alkyl" at R^1 , R^2 , R^3 and R^4 .

In the present invention, the "nonmetallic atoms necessary for forming a substituted or unsubstituted condensed benzo ring or condensed naphtho ring" means a bonding group necessary for forming a condensed benzo ring or condensed naphtho ring, which is a group of the formula



When the condensed benzo ring or condensed naphtho ring has a substituent, said bonding group may include a substituent.

Specific examples thereof include carbon atom, nitrogen atom, oxygen atom,
5 hydrogen atom, sulfur atom, halogen atom (e.g., fluorine atom, chlorine atom, bromine atom and iodine atom) and the like.


The substituent of the condensed benzo ring and condensed naphtho ring formed by the nonmetallic atoms at Z^1 and Z^2 is exemplified by sulfonic acid
10 group, carboxyl, hydroxy, halogen atom (e.g., fluorine atom, chlorine atom, bromine atom and iodine atom), cyano, substituted amino (e.g., dimethylamino, diethylamino, ethyl 4-sulfobutylamino, di-(3-sulfopropyl)amino and the like), and substituted or unsubstituted alkyl as defined above, which is bonded to the ring directly or via a divalent bonding group. Preferable divalent bonding group may
15 be, for example, -O-, -NHCO-, -NH SO_2 -, -NHCOO-, -NHCONH-, -COO-, -CO-, SO_2 -, and the like. The alkyl of substituted or unsubstituted alkyl that is bonded to the ring directly or via a divalent connection group is exemplified preferably by methyl, ethyl, propyl and butyl, and the substituent is preferably exemplified by sulfonic acid group, carboxyl and hydroxy.

20

The substituent of methine at $L^1 - L^7$ is exemplified by substituted or unsubstituted alkyl (defined above), halogen atom (defined above), substituted or unsubstituted aryl, lower alkoxy and the like. The aryl of "substituted or unsubstituted aryl" is exemplified by phenyl, naphthyl and the like, preferably
25 phenyl. Examples of the substituent include halogen atom (defined above, preferably chlorine atom) and the like. The substituted aryl includes, for example, 4-chlorophenyl and the like. The lower alkoxy is preferably linear or branched alkoxy having 1 to 6 carbon atoms, which is specifically methoxy, ethoxy, propoxy, butoxy, tert-butoxy, pentyloxy and the like, preferably methoxy
30 and ethoxy. In addition, the substituents of methine at $L^1 - L^7$ may be bonded each other to form a ring containing three methine groups, and this ring may further form a condensed ring with a ring containing different methine group. The ring containing three methine groups that is formed by the bonding of the

substituents of methine at $L^1 - L^7$ is exemplified by 4,4-dimethylcyclohexene ring and the like.

The conjugated methine chain consisting of the groups of $L^1 - L^7$, and having a
 5 ring is preferably the group of the formula (a):



10 wherein Z^3 denotes nonmetallic atoms necessary to form a 5- or 6-membered ring and A is hydrogen atom or a monovalent group.

The "nonmetallic atoms necessary to form a 5- or 6-membered ring" is
 15 exemplified by those mentioned above.

In the formula (a) and [III-2] to be mentioned later, - or 6-membered ring at Z^3 is exemplified by cyclopentene ring, cyclohexene ring, 4,4-dimethylcyclohexene ring and the like, with particular preference given to cyclopentene ring.

20

The monovalent group represented by A includes, for example, substituted or unsubstituted alkyl (defined above), substituted or unsubstituted aryl (defined above), substituted or unsubstituted aralkyl, lower alkoxy (defined above), substituted amino which is optionally substituted, alkylcarbonyloxy (e.g.,
 25 acetoxy), substituted or unsubstituted alkylthio, substituted or unsubstituted arylthio, cyano, nitro, halogen atom (defined above), and the like. As used herein, aralkyl of the "substituted or unsubstituted aralkyl" is exemplified by benzyl, 2-phenylethyl, 1-phenylethyl, 3-phenylpropyl and the like, and the substituent may be sulfonic acid group, carboxyl, hydroxy, substituted or
 30 unsubstituted alkyl (defined above), alkoxy (defined above), halogen atom (defined above), and the like. The substituted amino of the "substituted amino which is optionally substituted" includes, for example, alkylamino (e.g., methylamino, ethylamino and the like), dialkylamino (dimethylamino,

diethylamino and the like), diphenylamino, methylphenylamino, cyclic amino (e.g., morpholino, imidazolidino, ethoxycarbonylpiperadino and the like) and the like. The substituent with regard to the optional substitution of the "substituted amino which is optionally substituted" includes sulfonic acid group, carboxyl and the like. The alkylthio of the "substituted or unsubstituted alkylthio" may be, for example, methylthio, ethylthio and the like. Examples of the substituent include sulfonic acid group, carboxyl and the like. The arylthio of the "substituted or unsubstituted arylthio" is exemplified by phenylthio, naphthylthio and the like. Examples of the substituent include sulfonic acid group, carboxyl and the like.

10

The monovalent group represented by A is preferably fluorine atom, chlorine atom, dialkylamino (preferably having 6 or less carbon atoms, and optionally forming a ring) or morpholino. This group particularly preferably has a sulfonic acid group.

15

In the formula [I], r is preferably 1.

The pharmaceutically acceptable salt may be any as long as it forms a nontoxic salt with the compound of the formula [I]. Examples thereof include alkali metal salts such as sodium salt, potassium salt; salt of alkaline earth metal such as magnesium salt, calcium salt and the like; organic ammonium salt such as ammonium salt, triethyl ammonium salt, tributyl ammonium salt, pyridinium salt and the like; salt of amino acid such as lysine salt, arginine salt and the like. Particularly preferred is sodium salt causing less toxicity in the living body.

25

The fluorescent contrast agent to be used in a living body should be particularly water soluble. In the present invention, the near infrared fluorescent contrast agent has a noticeably improved water solubility by the introduction of 3 or more sulfonic acid groups into the above-mentioned compound. For superior water solubility, the number of the sulfonic acid groups is preferably 4 or more. For easy synthesis, the number of the sulfonic acid groups is not more than 10, preferably not more than 8. The improvement in the water solubility can be determined by measuring partition coefficient of each compound, which for example, may be measured in a two-phase system of butanol/water. More


30

specifically, introduction of 3 or more sulfonic acid groups results in a partition coefficient log Po/w of n-butanol/water of not more than -1.00.


The sulfonic acid groups are particularly preferably introduced into the positions of R¹, R², Z¹ and/or Z² of the formula [I] and R¹, R², R⁵, R⁷, R¹¹ and/or R¹³ of the formula [II].

In addition, these sulfonic acid groups are preferably introduced into L⁴ of the conjugated methine chain at the position A of the above-mentioned formula (a) via a divalent group such as alkylene.


Of the sodium salts of compounds of the formula [II] having three or more sulfonic acid groups in a molecule, preferred is a sodium salt of a compound wherein R¹ and R² are lower alkyl having 1 to 5 carbon atoms which is substituted by sulfonic acid group, and X and Y are the same or different and




each is a group of the formula wherein R¹⁷ and R¹⁸ are the same or different and each is unsubstituted lower alkyl having 1 to 5 carbon atoms alkyl, said salt having three or more sulfonic acid groups in a molecule, with particular preference given to a compound of the formula




Of the compounds of the formula [I] having three or more sulfonic acid groups in a molecule and pharmaceutically acceptable salts thereof, preferred is a sodium salt of a compound of the formula [III-1]




- 5 wherein $L^1 - L^7$ are as defined above, R^{19} and R^{20} are lower alkyl having 1 to 5 carbon atoms which is substituted by sulfonic acid group, and R^{21} to R^{28} are the same or different and each is hydrogen atom, sulfonic acid group, carboxyl group, hydroxyl group, alkyl(sulfoalkyl)amino group, bis(sulfoalkyl)amino group, sulfoalkoxy group, (sulfoalkyl)sulfonyl group or (sulfoalkyl)amino-sulfonyl group,
- 10 and X' and Y' are the same or different and each is a group of the formula




wherein R^{17} and R^{18} are as defined above, said salt having three or more sulfonic acid groups in a molecule, with particular preference given to the compound of the following formula.



Of the sodium salts of the compounds of the formula [III-1] having three or more sulfonic acid groups in a molecule, preferred is a sodium salt of a compound of the formula [III-2]



wherein R^{19} - R^{28} , X' and Y' are as defined above, Z^3 is nonmetallic atoms necessary to form a 5- or 6-membered ring and A is hydrogen atom or a monovalent group, said salt having three or more sulfonic acid groups in a molecule, with particular preference given to the compound of the following formula



The compounds contained in the inventive near infrared fluorescent contrast agent may be any as long as it has formula [I] or [II], as well as 3 or more, preferably 4 or more, sulfonic acid groups in a molecule. These compounds can be synthesized according to a known production method of cyanine dye compounds disclosed in *The Cyanine Dyes and Related Compounds*, F.M. Hamer, John Wiley and Sons, New York, 1964, *Cytometry*, 10, 3-10 (1989), *Cytometry*, 11, 418-430 (1990), *Cytometry*, 12, 723-730 (1990), *Bioconjugate Chem.* 4, 105-111 (1993), *Anal. Biochem.*, 217, 197-204 (1994), *Tetrahedron*, 45, 4845-4866 (1989), EP-A-0591820A1, EP-A-0580145A1, and the like. Alternatively, they can be semisynthesized from a commercially available cyanine dye compound by a known method. To be specific, they can be synthesized by reacting a dianyl compound and a heterocyclic quaternary salt.


The compound of the formula [I] of the present invention can be synthesized by, for example, the following method.

(i) when $r=0$


5 (a) $L^1=L^5$, $X=Y$, $R^1=R^2$ and $Z^1=Z^2$

A hetero ring quaternary salt compound (2 moles) of the formula


[IV-1]



wherein L^1 , X , Z^1 and R^1 are as defined above, and a dianyl compound (1 mole) of the formula [V-1]



wherein L^2 , L^3 and L^4 are as defined above, are reacted in the presence of a base and a solvent to give a compound of the formula [VI-1]




wherein L^1 , L^2 , L^3 , L^4 , R^1 , Z^1 and X are as defined above, and this compound [VI-1] (1 mole) and a necessary molar amount of the compound of the formula [VII]

30 $T^1 - Na$ [VII]


wherein T^1 is an organic acid residue, are reacted to give a sodium salt of the compound of the above-mentioned formula [VI-1].

(b) $L^1 \neq L^5$ or $X \neq Y$ or $R^1 \neq R^2$ or $Z^1 \neq Z^2$


A hetero ring quaternary salt compound (1 mole) of the above-mentioned formula [IV-1] and the above-mentioned diaryl compound (1 mole) of the formula [V-1] are reacted in the presence of a base and a solvent to give a compound of the formula [VIII-1]



10 wherein L^1 , L^2 , L^3 , L^4 , R^1 , Z^1 and X are as defined above, and this compound [VIII-1] (1 mole) and a hetero ring quaternary salt compound (1 mole) of the formula [XI-1]



wherein L^5 , Y , Z^2 and R^2 are as defined above, are reacted to give a compound of the formula [X-1]




25 wherein L^1 , L^2 , L^3 , L^4 , L^5 , R^1 , R^2 , Z^1 , Z^2 , X and Y are as defined above, and this compound of the formula [X-1] (1 mole) and a necessary molar amount of the above-mentioned compound of the formula [VII] are reacted to give a sodium salt of the compound of the above-mentioned formula [X-1].

(ii) when $r=1$


(a) $L^1=L^7$, $X=Y$, $R^1=R^2$ and $Z^1=Z^2$

A hetero ring quaternary salt compound (2 moles) of the formula [IV-1]




5

wherein L^1 , X , Z^1 and R^1 are as defined above, and a dianyl compound (1 mole) of the formula [V-2]



10

wherein L^2 , L^3 , L^4 , L^5 and L^6 are as defined above, are reacted in the presence of a base and a solvent to give a compound of the formula [VI-2]



15

wherein L^1 , L^2 , L^3 , L^4 , L^5 , L^6 , R^1 , Z^1 and X are as defined above, and this compound [VI-2] (1 mole) and a necessary molar amount of the compound of the formula [VII]




wherein T^1 is as defined above, are reacted to give a sodium salt of the compound of the above-mentioned formula [VI-2].


(b) $L^1 \neq L^7$ or $X \neq Y$ or $R^1 \neq R^2$ or $Z^1 \neq Z^2$

A hetero ring quaternary salt compound (1 mole) of the above-mentioned formula [IV-1] and the above-mentioned dianyl compound (1 mole) of the formula [V-2] are reacted in the presence of a base and a solvent to give a compound of the formula [VIII-2]


30



wherein L^1 , L^2 , L^3 , L^4 , L^5 , L^6 , R^1 , Z^1 and X are as defined above, and this compound [VIII-2] (1 mole) and a hetero ring quaternary salt compound (1 mole) of the formula [IX-2]



15 wherein L^7 , Y , Z^2 and R^2 are as defined above, are reacted to give a compound of the formula [X-2]



25 wherein L^1 , L^2 , L^3 , L^4 , L^5 , L^6 , L^7 , R^1 , R^2 , Z^1 , Z^2 , X and Y are as defined above, and this compound of the formula [X-2] (1 mole) and a necessary molar amount of the above-mentioned compound of the formula [VII] are reacted to give a sodium salt of the compound of the above-mentioned formula [X-2].


(iii) when $r=2$

When r is 2, L^6 and L^7 overlap in the formula [I]. To avoid this, the overlapping L^6 and L^7 are referred to as L^8 and L^9 for clarification.


30 (a) $L^1=L^9$, $X=Y$, $R^1=R^2$ and $Z^1=Z^2$

A hetero ring quaternary salt compound (2 moles) of the formula


[IV-1]



wherein L^1 , X , Z^1 and R^1 are as defined above, and a dianyl compound (1 mole) of the formula [V-3]



wherein L^2 , L^3 , L^4 , L^5 , L^6 and L^7 are as defined above and L^8 is an optionally substituted methine group, are reacted in the presence of a base and a solvent to give a compound of the formula [VI-3]



wherein L^1 , L^2 , L^3 , L^4 , L^5 , L^6 , L^7 , L^8 , R^1 , Z^1 and X are as defined above, and this compound [VI-3] (1 mole) and a necessary molar amount of the compound of the formula [VII]




wherein T^1 is as defined above, are reacted to give a sodium salt of the compound of the above-mentioned formula [VI-3].

(b) $L^1 \neq L^9$ or $X \neq Y$ or $R^1 \neq R^2$ or $Z^1 \neq Z^2$


A hetero ring quaternary salt compound (1 mole) of the above-mentioned formula [IV-1] and the above-mentioned dianyl compound (1 mole) of the formula [V-3] are reacted in the presence of a base and a solvent to give a

compound of the formula [VIII-3]




wherein L^1 , L^2 , L^3 , L^4 , L^5 , L^6 , L^7 , L^8 , R^1 , Z^1 and X are as defined above, and this compound [VIII-3] (1 mole) and a hetero ring quaternary salt compound (1 mole) of the formula [IX-3]

10



wherein Y , Z^2 and R^2 are as defined above and L^9 is an optionally substituted methine group, are reacted to give a compound of the formula [X-3]



25 wherein L^1 , L^2 , L^3 , L^4 , L^5 , L^6 , L^7 , L^8 , L^9 , R^1 , R^2 , Z^1 , Z^2 , X and Y are as defined above, and this compound of the formula [X-3] (1 mole) and a necessary molar amount of the above-mentioned compound of the formula [VII] are reacted to give a sodium salt of the compound of the above-mentioned formula [X-3].

30 The necessary molar amount of the compound of the formula [VII] is not less than the amount equivalent to the amount of sodium contained in one molecule of the objective sodium salt of the compound of the formula [I].

The substituent of the substituted methine group at L⁸ and L⁹ is exemplified by those mentioned with regard to the substituent of the above-mentioned methine groups at L¹ to L⁷.

- 5 In the synthetic methods of the above-mentioned (i), (ii) and (iii), the reaction of the compounds [IV-1] and [V-1], that of the compounds [VIII-1] and [XI-1], that of the compounds [IV-1] and [V-2], that of the compounds [VIII-2] and [IX-2], that of the compounds [IV-1] and [V-3] and that of the compounds [VIII-3] and [IX-3] proceed at a temperature of -20 - 80 , preferably -10 - 40 , preferably in
10 the presence of an acylating agent such as acetic anhydride.


- In the synthetic methods of the above-mentioned (i), (ii) and (iii), the reaction of the compounds [IV-1] and [VII], that of the compounds [X-1] and [VII], that of the compounds [VI-2] and [VII], that of the compounds [X-2] and [VII], that of the
15 compounds [VI-3] and [VII] and that of the compounds [X-3] and [VII] proceed at a temperature of preferably 0 - 40 , preferably in the presence of a solvent such as alcohol and water.


- In the synthetic methods of the above-mentioned (i), (ii) and (iii), the base to be
20 used may be, for example, triethylamine, tributylamine, pyridine, diazabicycloundecene, sodium methoxide and the like; the solvent to be used may be, for example, an amide compound such as N,N-dimethylacetamide, N-methylpyrrolidone and N,N-diethylformamide or alcohols such as methanol; and the organic acid residue may be, for example, CH₃COO and the like.

- 25 With regard to the production of various pharmaceutically acceptable salts of the compounds of the aforementioned formula [I], ammonium salt and potassium salt of the compounds of the formula [I] can be obtained by, for example, substituting the compound of the formula [VII] used in the above-mentioned
30 synthetic methods (i), (ii) and (iii) with a compound of the formula [VII] wherein the sodium atom has been changed to ammonium group or potassium atom; and different cationic salts of the compounds of the aforementioned formula [I] can be obtained by converting said ammonium salt and potassium salt to


different cationic salts by the use of ion exchange resins as necessary.

The compound of the above-mentioned formula [I] inclusive of the compound of the formula [II] to be used in the present invention are specifically exemplified in the following, to which the present invention is not limited.







(11)




(12)




(13)




(14)




(15)




(16)




(17)




(18)




(19)




(20)




(21)




(22)




(23)




(24)




(25)




(26)







(27)




(28)







(33)




(34)




(35)




(36)




(37)




(38)




(39)




(40)




(41)




(42)




(43)




(44)




(45)




(46)




(47)




(48)




(49)




(50)




(51)




(52)



(53)



(54)



The above-mentioned compound to be contained in the near infrared fluorescent contrast agent of the present invention shows absorbance and fluorescence in the near infrared light region of 700 – 1300 nm, particularly about 700 – 900 nm, and has a molar absorption coefficient of not less than 100,000.

The near infrared fluorescent contrast agent of the present invention is subject to no particular limitation as long as it contains a compound of the formula [I] or formula [II] and/or a pharmaceutically acceptable salt thereof, and has 3 or more, preferably 4 or more, sulfonic acid groups in a molecule. This compound or a salt thereof alone or in combination may be contained in said contrast agent.

To be specific, said contrast agent includes said compound or said compound suspended or dissolved in a solvent, such as injectable distilled water, physiological saline, Ringer solution and the like. Where necessary, pharmacologically acceptable additives such as carrier, excipient and the like

may be added. These additives contain substances such as pharmacologically acceptable electrolyte, buffer, detergent and a substance for adjusting osmotic pressure and improving stability and solubility (e.g., cyclodextrin, liposome and the like). Various additives generally used in the pertinent fields may be used. The near infrared fluorescent contrast agent of the present invention is preferably produced through a sterilization process when it is intended for pharmaceutical use.

Said contrast agent can be administered to a living body by injecting, spraying or coating, intravascularly (venous, arterial), orally, intraperitoneally, percutaneously, subcutaneously, intracystically or intrabronchially. Preferably, the agent is administered into blood vessels in the form of an aqueous agent, emulsion or suspension.

The dose of the near infrared fluorescent contrast agent of the present invention is not particularly limited insofar as the dose enables detection of the site to be ultimately diagnosed. It is appropriately adjusted depending on the kind of compound to be used that emits near infrared fluorescence, age, body weight and target organ of administration subjects, and the like. Typically, the dose is 0.1 – 100 mg/kg body weight, preferably 0.5 – 20 mg/kg body weight, in the amount of said compound.

The contrast agent of the present invention can be appropriately used for various animals other than human. The administration form, route and dose are suitably determined depending on the body weight and conditions of the target animals.

In the present invention, moreover, the above-mentioned compound of the formula [I], particularly preferably [II], having 3 or more, preferably 4 or more, sulfonic acid groups in a molecule tends to be noticeably accumulated in tumor tissues. Utilizing this characteristic, a tumor tissue can be specifically imaged using the inventive fluorescent contrast agent. In addition, a series of said

compounds can reside in blood vessel for a long time and they are expected to serve well as angiography contrast agents.

The fluorescence imaging method of the present invention is characterized by the use of the inventive near infrared fluorescent contrast agent. This method is practiced following known methods, and each parameter, such as excitation wavelength and fluorescence wavelength to be detected, is appropriately determined to achieve optimal imaging and evaluation, depending on the kind of near infrared fluorescent contrast agent to be administered and administration targets. The time spent from administration of the inventive near infrared fluorescent contrast agent to determination target to the initiation of determination by the inventive fluorescence imaging method varies depending on the kind of the near infrared fluorescent contrast agent to be used and administration targets. For example, when the agent contains a compound of the formula [I] for tumor imaging, the lapse time will be about 4-120 hours after administration. In the case of compound of formula [II], the lapse time will be about 24-120 hours after administration. When the lapse time is too short, the fluorescence is so intense that the target site and other site cannot be clearly divided. When it is too long, said contrast agent may be cleared from the body. When imaging of blood vessel is desired, the compound of the formula [I] or formula [II] is detected immediately after administration or in about 30 minutes thereafter.

The method typically includes the following steps.

That is, a near infrared fluorescent contrast agent of the present invention is administered to a detection target and the detection target is exposed to an excitation light from an excitation light source. Then, fluorescence from the near infrared fluorescent contrast agent, which has been caused by said excitation light, is detected with a fluorescence detector.

The wavelength for excitation varies depending on the near infrared fluorescent contrast agent to be used. It is free of limitation as long as said

09767394-051604

compound efficiently emits fluorescence in the near infrared region. Preferably, a near infrared light having superior biotransmission capability is used.

The wavelength of the near infrared fluorescence to be detected also varies depending on the contrast agent to be used. In general terms, an excitation light having a wavelength of 600-1000 nm, preferably 700-850 nm, is used and near infrared fluorescence in a region at a wavelength of 700-1000 nm, preferably 750-900 nm, is detected. In this case, the excitation light source may be a conventional excitation light source, such as various lasers (e.g., ion laser, dye laser and semiconductor laser), halogen light source, xenon light source and the like. Where necessary, various optical filters may be used to obtain optimal excitation wavelength. Likewise, fluorescence may be detected using various optical filters to pick up only fluorescence from said near infrared fluorescent contrast agent.

The detected fluorescence is data-processed as fluorescence information and used to generate fluorescence images that can be recorded. The fluorescence images are generated by irradiating a wide area including the target tissue, detecting fluorescence with a CCD camera and image-processing the obtained fluorescence information. Alternatively, an optical CT device may be used, an endoscope may be used, or a fundus camera may be used.

The fluorescence imaging method of the present invention enables visualizing systemic diseases, tumors, blood vessels and the like without damaging a living body.




The present invention is explained in more detail by way of Examples and Experimental Examples, to which the present invention is not limited. The compound numbers in the following Examples and Experimental Examples correspond to those of the compounds explained by structural formulas.

The compound wherein a symbol designating "potassium salt", "calcium salt" or "pyridinium salt" is indicated after the compound number (e.g. compound (29) K salt) means a compound that is the same as the compound expressed by the compound number (sodium salt) except that the counter ion is potassium salt, calcium salt or pyridinium salt instead of sodium salt. For example, "compound (31)K salt" means a compound that is the same as the compound (31) except that the counter ion is potassium instead of sodium; "compound (31)Ca salt" means a compound that is the same as compound (31) except that the counter ion is calcium instead of sodium; and "compound (31)pyridinium" salt means a compound that is the same as compound (31) except that the counter ion is pyridinium instead of sodium.

The synthetic method of the compound to be contained in the near infrared fluorescent contrast agent of the present invention as an active ingredient is explained in Examples.

The following synthetic methods mostly consist of reactions of heterocyclic quaternary salt compound shown in Table 1 and dianyl compounds shown in Tables 2 and 3.

Table 1 Heterocyclic quaternary salt compound

Symbol	Structural formula [Registration No. in Chemical Abstract (CA)]	Source/Synthetic method
Q1	 <p>(138913-76-5)</p>	In the same manner as Q2, corresponding indolenin derivative is reacted with butanesultone.
Q2	 <p>(113995-56-5)</p>	JP-A 63-55544 EP 251282
Q3	 <p>(76588-81-3)</p>	JP-A 2-233658 CA 114:122053






Q4	 (183272-36-8)	In the same manner as Q3, corresponding indolenin derivative is reacted with 2- bromoethanesulfonic acid.
----	--	--

Table 2 Dianyl compound-1

Symbol	Structural formula [Registration No. in Chemical Abstract (CA)]	Source/Synthetic method
A1	 (1497-49-0) • HCl	Reagent commercially available from Aldrich and others
A2	 (1979-58-4) • HCl	JP-A 8-295658 CA 126:90721
A3	 (53019-66-2) • HCl	Zh.Org.Khim., 13(6) 1189-92 (1977) CA 87:102034
A4	 (63856-99-5) • HCl	Zh.Org.Khim., 13(6) 1189-92 (1977) CA 87:102034

Table 3 Dianyl compound-2

Symbol	Structural formula [Registration No. in Chemical Abstract (CA)]	Source/Synthetic method
A5	 <p>(77146-76-0) • HCl</p>	Nukleofil' nye Reacts. Karbonil' nykn Soedin (1982), 52-53 CA 101:130179
A6	 <p>(125577-71-1) • HCl</p>	Ger Offem, DE 2928184 CA 94:176696
A7	 <p>(56709-94-5) • HCl</p>	Zh.Org.Khim., 18(10) 2176-9 (1982) CA 98:73808
A8	 <p>(63857-00-1) • HCl</p>	Zh.Org.Khim., 13(6) 1189-92 (1977) CA 87:102034

Examples

In the following Examples, the compounds are referred to with the symbols (e.g., A1, Q1 and the like) used in Tables 1 to 3 for the convenience's sake.

Example 1 : Synthesis of compound (29)

To heterocyclic quaternary salt compound Q1 (5 g) were added methanol (100 ml), N,N-dimethylformamide (25 ml), triethylamine (5.6 ml), dianyl compound A1 (1.83 g) and acetic anhydride (3 ml), and the mixture was stirred at room temperature for 4 hours. Triethyl amine (2.2 ml) and acetic anhydride (2 ml) were added, and the mixture was stirred at room temperature for 3 hours. The insoluble matter was filtered off, and a solution of sodium acetate (2 g) in methanol (15 ml) was added to the filtrate, which was followed by stirring at room temperature for 1 hour. The resulting crystals were collected by filtration and washed with a small amount of methanol. To the obtained crude crystals (3.5 g) was added water (20 ml) for dissolution. Sodium acetate (1 g) was added, and then methanol (30 ml) was added, which was followed by stirring for 1 hour. The resulting crystals were collected by filtration, washed with a small amount of methanol and dried to give 3 g of compound (29). The obtained compound (29) showed yellow in a flame test.

Maximum wavelength of absorbance (H₂O) : 780 nm

Molar absorption coefficient (H₂O) : 243,000

Maximum wavelength of fluorescence emission (H₂O) : 802 nm

The infrared absorption spectrum was measured for the obtained compound (29) by potassium bromide tablet method using a Fourier transform infrared spectrometer (VALOR-III, manufactured by JASCO). The following peaks were detected. The spectrum is shown in Fig. 11.

IR (<max(KBr)) : 1414, 1086, 1037, 995, 889 cm⁻¹

Example 2 : Synthesis of compound (34)

To heterocyclic quaternary salt compound Q2 (2.13 g) was added methanol (20 ml) and the mixture was cooled to 10 . Thereto were added dianyl

compound A2 (0.75 g), triethylamine (4 ml) and acetic anhydride (2 ml), and the mixture was stirred for 20 minutes. Acetic anhydride (2 ml) was added, and the mixture was stirred at 10 °C for 4 hours. The insoluble matter was filtered off, and a solution of sodium acetate (2 g) in a small amount of methanol was added to the filtrate. The resulting crystals were collected by filtration and washed with a small amount of methanol. To the obtained crude crystals was added water (7ml) for dissolution. Methanol (7 ml) was added to precipitate crystals. The resulting crystals were collected by filtration, washed with a small amount of methanol and dried to give 1.2 g of compound (34). The obtained compound (34) showed yellow in a flame test.

Maximum wavelength of absorbance (H₂O) : 794 nm

Molar absorption coefficient (H₂O) : 176,000

Maximum wavelength of fluorescence emission (H₂O) : 812 nm

Example 3 : Synthesis of compound (6)

To heterocyclic quaternary salt compound Q3 (9.5 g) are added methanol (50 ml), triethylamine (7 ml), dianyl compound A3 (3.1 g) and acetic anhydride (3.9 ml), and the mixture is stirred at room temperature for 7 hours. The insoluble matter is filtered off, and a solution of sodium acetate (5 g) in a small amount of methanol is added to the filtrate. The mixture is stood overnight. The resulting crystals are collected by filtration and washed with a small amount of methanol. To the crystals is added water (30 ml) for dissolution. Sodium acetate (2 g) is added, and then methanol (30 ml) is added. The resulting crystals are collected by filtration, washed with a small amount of methanol and dried to give compound (6).

Example 4 : Synthesis of compound (45)

To heterocyclic quaternary salt compound Q3 (4.8 g) were added methanol (50 ml), triethylamine (4 ml), dianyl compound A4 (1.7 g) and acetic anhydride (2 ml), and the mixture was stirred at room temperature for 3 hours. The insoluble matter was filtered off, and a solution of sodium acetate (4 g) in a small amount of methanol was added to the filtrate. The resulting crystals

were collected by filtration and washed with a small amount of methanol. To the crystals was added water (10 ml) for dissolution. Then methanol (10 ml) was added. The resulting crystals were collected by filtration, washed with a small amount of methanol and air dried to give 1.6 g of a compound that is the same as compound (45) except that the substituent on the methine chain is -Cl instead of -SCH₂CH₂SO₃ Na.

The above step was repeated to give 4.2 g of said compound. Thereto were added water (30 ml), triethylamine (1.2 ml) and sodium 2-mercaptoethanesulfonate (0.8 g), and the mixture was stirred at room temperature for 4 hours. The insoluble matter was filtered off, and a solution of sodium acetate (2 g) in a small amount of water was added to the filtrate. The resulting crystals were collected by filtration, washed with methanol (20 ml) and air dried to give 2.3 g of compound (45). The obtained compound (45) showed yellow in a flame test.

Maximum wavelength of absorbance (H₂O) : 815 nm

Molar absorption coefficient (H₂O) : 196,000

Maximum wavelength of fluorescence emission (H₂O) : 827 nm

Example 5 : Synthesis of compound (2)

To heterocyclic quaternary salt compound Q3 (4.7 g) are added methanol (25 ml), triethylamine (2.8 ml), dianyl compound A5 (1.5 g) and acetic anhydride (2.4 ml), and the mixture is stirred at room temperature for 1 hour. Thereto are further added triethyl amine (3.5 ml) and acetic anhydride (1.5 ml), and the mixture is stirred at room temperature for 3.5 hours. The insoluble matter is filtered off, and a solution of sodium acetate (3 g) in a small amount of methanol is added to the filtrate. The mixture is stirred at room temperature for 1 hour. The resulting crystals are collected by filtration and washed with a small amount of methanol. To the crystals is added water (15 ml) for dissolution. Then methanol (15 ml) is added. The resulting crystals are collected by filtration, washed with a small amount of methanol and dried to give compound (2).

Example 6 : Synthesis of compound (43)

To heterocyclic quaternary salt compound Q3 (3.75 g) were added methanol (25 ml), triethylamine (3.5 ml), dianyl compound A6 (1.95 g) and acetic anhydride (2.4 ml), and the mixture was stirred at room temperature for 1 hour. The insoluble matter was filtered off, and a solution of sodium acetate (3.9 g) in a small amount of methanol was added to the filtrate. The mixture was stirred at room temperature for 1 hour. The resulting crystals were collected by filtration and washed with a small amount of methanol. To the crystals was added water (10 ml) for dissolution. Sodium acetate (2 g) was added, and then methanol (10 ml) was added. The resulting crystals were collected by filtration, washed with a small amount of methanol and dried to give 1.8 g of compound (43). The obtained compound (43) showed yellow in a flame test.

Maximum wavelength of absorbance (H₂O) : 773 nm

Molar absorption coefficient (H₂O) : 204,000

Maximum wavelength of fluorescence emission (H₂O) : 789 nm

Example 7 : Synthesis of compound (4)

To heterocyclic quaternary salt compound Q3 (3.5 g) are added methanol (20 ml), triethylamine (3.5 ml), dianyl compound A7 (1.2 g) and acetic anhydride (1.9 ml), and the mixture is stirred at room temperature for 10 hours, and then stood overnight. The mixture is stirred under heating at 50 °C for 5 hours. Water (2 ml) is added and the insoluble matter is filtered off. A solution of sodium acetate (5 g) in a small amount of water is added to the filtrate. The mixture is stirred at room temperature for 30 minutes. The resulting crystals are collected by filtration and washed with a small amount of methanol and dried to give compound (4).

Example 8 : Synthesis of compound (31)

To heterocyclic quaternary salt compound Q4 (3.5 g) were added methanol (35 ml), triethylamine (3.5 ml) and acetic anhydride (2 ml), and dianyl compound A2 (1.8 g) was added portionwise with stirring. The mixture was further stirred for 1 hour. Acetic anhydride (2 ml) was added, and the mixture was stirred at

room temperature for 5 hours. The insoluble matter was filtered off, and a solution of sodium acetate (4 g) in a small amount of methanol was added to the filtrate. The resulting crystals were collected by filtration and washed with a small amount of methanol. To the crystals was added water (10 ml) for dissolution. Then methanol (10 ml) was added, and the mixture was stirred at room temperature for 2 hours. The resulting crystals were collected by filtration, washed with a small amount of methanol and dried to give 1.3 g of compound (31). The obtained compound (31) showed yellow in a flame test.

Maximum wavelength of absorbance (H₂O) : 755 nm

Molar absorption coefficient (H₂O) : 228,000

Maximum wavelength of fluorescence emission (H₂O) : 774 nm

The infrared absorption spectrum was measured for the obtained compound (31) by potassium bromide tablet method using a Fourier transform infrared spectrometer (VALOR-III, manufactured by JASCO). The following peaks were detected. The spectrum is shown in Fig. 12.

IR (ν_{max} (KBr)) : 1518, 1183, 1149, 1111, 995 cm⁻¹

Example 9 : Synthesis of compound (41)

To heterocyclic quaternary salt compound Q1 (12 g) were added methanol (120 ml), triethylamine (13.6 ml), dianyl compound A8 (4.4 g) and acetic anhydride (2.4 ml), and the mixture was stirred for 30 minutes. Acetic anhydride (2.4 ml) was added and the mixture was stirred for 1.5 hours, and then acetic anhydride (2.4 ml) was added and the mixture was stirred at room temperature for 6 hours. Heterocyclic quaternary salt compound Q1 (1 g), triethyl amine (3 ml) and acetic anhydride (3 ml) were further added and the mixture was stirred at room temperature for 2 hours. The mixture was stood overnight. Sodium acetate (5 g) was added and the resulting crystals were collected by filtration and washed with a small amount of methanol. To the obtained crude crystals was added water (200 ml). The insoluble matter was filtered off, and sodium acetate (10 g) was added to the filtrate. The resulting crystals were collected by filtration and washed with a small amount of

methanol. To the crystals were added water (200 ml) and triethylamine (10 ml), and a solution of sodium acetate (10 g) in methanol (100 ml) was added to give crystals. This step was repeated twice. The resulting crystals were collected by filtration, washed with a small amount of methanol and dried to give 9.7 g of compound (41). The obtained compound (41) showed yellow in a flame test.

Maximum wavelength of absorbance (H_2O) : 811 nm

Molar absorption coefficient (H_2O) : 230,000

Maximum wavelength of fluorescence emission (H_2O) : 822 nm

Example 10 : Synthesis of compound (3)

According to Example 5, heterocyclic quaternary salt compound Q3 and the corresponding dianyl compound are used to give compound (3).

Example 11

In the same manner as in the synthesis of compound (29) in Example 1 except that potassium acetate (2 g) was used instead of sodium acetate (2 g), a compound that is the same as compound (29) except that the counter ion was potassium instead of sodium was obtained. Hereinafter this compound is referred to as compound (29) K salt. The obtained compound (29) K salt showed purple in a flame test.

Maximum wavelength of absorbance (H_2O) : 780 nm

Molar absorption coefficient (H_2O) : 254,000

Maximum wavelength of fluorescence emission (H_2O) : 800 nm

Other aforementioned compounds are treated in the same manner as in this Example to give compounds having potassium counter ion instead of sodium.

These compounds having potassium counter ion are distinguished from the above compounds by attaching "K salt" after the corresponding compound numbers.

Example 12

In the same manner as in Example 11, compound (6) K salt was obtained. The obtained compound (6) K salt showed purple in a flame test.

Maximum wavelength of absorbance (H₂O) : 788 nm

Molar absorption coefficient (H₂O) : 226,000

Maximum wavelength of fluorescence emission (H₂O) : 806 nm

Example 13

In the same manner as in Example 11, compound (2) K salt was obtained. The obtained compound (2) K salt showed purple in a flame test.

Maximum wavelength of absorbance (H₂O) : 743 nm

Molar absorption coefficient (H₂O) : 266,000

Maximum wavelength of fluorescence emission (H₂O) : 762 nm

Example 14

In the same manner as in Example 11, compound (4) K salt was obtained. The obtained compound (4) K salt showed purple in a flame test.

Maximum wavelength of absorbance (H₂O) : 753 nm

Molar absorption coefficient (H₂O) : 212,000

Maximum wavelength of fluorescence emission (H₂O) : 767 nm

Example 15

In the same manner as in Example 11, compound (3) K salt was obtained. The obtained compound (3) K salt showed purple in a flame test.

Maximum wavelength of absorbance (H₂O) : 751 nm

Molar absorption coefficient (H₂O) : 241,000

Maximum wavelength of fluorescence emission (H₂O) : 767 nm

Example 16

The compound (6) K salt (50 mg) was dissolved in a small amount of water and passed through an ion exchange resin to convert potassium of the compound (6) K salt to proton. Thereto was added methanol saturated with

sodium acetate to allow precipitation of crystals. This procedure was repeated twice. The resulting crystals were collected by filtration, washed with a small amount of methanol and dried to give (32 mg) of compound (6). The obtained compound (6) showed yellow in a flame test.

The infrared absorption spectrum was measured for the obtained compound (6) by potassium bromide tablet method using a Fourier transform infrared spectrometer (VALOR-III, manufactured by JASCO). The following peaks were detected. The spectrum is shown in Fig. 13.

IR (ν_{max} (KBr)) : 1395, 1372, 1188, 1102, 1020 cm^{-1}

Example 17 : Synthesis of compound (54)

To heterocyclic quaternary salt compound Q4 (3.5 g) were added methanol (20 ml), triethylamine (3.5 ml) and acetic anhydride (2 ml), and dianyl compound A1 (1.4 g) was added portionwise with stirring. The mixture was further stirred for 20 minutes. Acetic anhydride (1 ml) was added, and the mixture was stirred at room temperature for 1.5 hours. The insoluble matter was filtered off, and a solution of sodium acetate (4 g) in a small amount of methanol was added to the filtrate. The resulting crystals were collected by filtration and washed with a small amount of methanol. The crystals were dissolved in a small amount of water. Then the solution was diluted with methanol (10 ml), and the mixture was stirred at room temperature for 1 hour. The resulting crystals were collected by filtration, washed with a small amount of methanol and dried to give 1.5 g of compound (54). The obtained compound (54) showed yellow in a flame test.

Maximum wavelength of absorbance (H_2O) : 743 nm

Molar absorption coefficient (H_2O) : 244,000

Maximum wavelength of fluorescence emission (H_2O) : 766 nm

The infrared absorption spectrum was measured for the obtained compound (54) by potassium bromide tablet method using a Fourier transform infrared spectrometer (VALOR-III, manufactured by JASCO). The following peaks were

detected. The spectrum is shown in Fig. 14.



IR (ν_{max} (KBr)) : 1511, 1421, 1099, 1004, 926 cm^{-1}

Experimental Example 1

The partition coefficient ($\log P_{\text{o/w}}$) of n-butanol/water was determined with respect to compound (29), compound (43), compound (45), compound (31), compound (3) K salt, compound (11) [available from Nippon Kankoh-Shikiso Kenkyusho CO., LTD. as NK-3261], compound (6) K salt, compound (2) K salt, compound (4) K salt, compound (34) and compound (54).

As a control compound, used were NK-1967 (Nippon Kankoh-Shikiso Kenkyusho CO., LTD.) and ICG (Tokyo Kasei Kogyo) having only 2 sulfonic acid groups in a molecule. The results are shown in Table 4.

Table 4

Compound	Number of sulfonic acid group	log Po/w (butanol/water)
Compound (29)	6	-2.00 or less
Compound (43)	5	-2.00 or less
Compound (45)	5	-2.00 or less
Compound (31)	4	-2.00 or less
Compound (3) K salt	4	-2.00 or less
Compound (11) (NK-3261)	4	-2.00 or less
Compound (6) K salt	4	-2.00 or less
Compound (2) K salt	4	-2.00 or less
Compound (4) K salt	4	-1.51
Compound (34)	4	-1.49
Compound (54)	4	-2.00 or less
NK-1967 	2	0.34
ICG 	2	1.41

Experimental Example 2 : Fluorescence imaging test (1)

Tumor tissue pieces of mouse colon carcinoma (colon 26 carcinoma) were subcutaneously grafted to the left breast of BALB/c nude mice (5 weeks old, Clea Japan, Inc.). Ten days later when the tumor grew to a diameter of about 8 mm, the mice were subjected to the test.

As a fluorescence excitation light source, a titanium sapphire laser was used. The test mice were uniformly exposed to the laser light using a ring type light guide (Sumita Optical Glass Co.) wherein dispersion of irradiation was within 10%. The irradiation power output was adjusted so that it was about 40 $\mu\text{W}/\text{cm}^2$ near skin surface of the mice. The fluorescence was excited at the maximum excitation wavelength of each compound and fluorescence emission from the mice was detected and photographed through a short wavelength cutoff filter (IR84, IR86, IR88, Fuji Photo Film CO., LTD.) with a CCD camera (C4880, Hamamatsu Photonics K.K.). The cutoff filter was selected to fit the excitation wavelength of the compound. The exposure time was adjusted depending on the fluorescence intensity of each compound.

The test compounds used were compound (29), compound (31) and compound (6) K salt of the present invention, and NK-1967 and ICG having only 2 sulfonic acid groups in a molecule as control compounds. Each test compound (0.5 mg/ml) was dissolved in distilled water and administered to the mice via a tail vein. The dose was 5.0 mg/kg for compound (31), compound (6) K salt, NK-1967 and ICG, and 0.5 mg/kg for compound (29). At 24 hours after the administration of the compounds, the mice were anesthetized with diethyl ether and fluorescent light images of the entire body of the mice was photographed. The results are shown in Figs. 1 to 5.

The compound (29) having a benzotricarbocyanine structure and six sulfonic acid groups, as well as compound (6) K salt and compound (31) both having a tricarbocyanine structure and four sulfonic acid groups obviously generated clearer images of tumor as compared to control compounds (NK-1967 having

benzotricarbocyanine structure and ICG having tricarbocyanine structure) having two sulfonic acid groups. In particular, compound (29) could clearly depict the tumor even at a low dose and was noticeably effective.

Experimental Example 3 : Fluorescence imaging test (2)

Nude mice were used for the test. compound (29) of the present invention and control compound ICG were intravenously injected from the tail vein at a dose of 5.0 mg/kg each under sevoflurane continuous inhalation anesthesia. At the same time, intermittent photographing of the fluorescence images was initiated. For photographing of the fluorescence images, exposure to excitation laser beam and extraction of fluorescence through a filter were done, wherein exposure time was one second. At 20 seconds after the administration of the compounds, blood vessel was suitably imaged. The fluorescence images were photographed until 5 minutes after the administration. Figs. 6 to 9 show fluorescence images of the entire body of the mice at 20 seconds and 5 minutes after the administration.

ICG failed to contrastively show blood vessel in 5 minutes, whereas compound (29) could image the blood vessel for longer time than ICG.

Experimental Example 4 : Residence in blood vessel

In the same manner as in Experimental Example 2, tumor tissue pieces were grafted to CDF₁ mice (female, 5 weeks old, Japan SLC, Inc.), and about 2 weeks later when the tumor grew to a diameter of about 1 cm, the mice were subjected to the test.

The test compounds were compound (29) K salt and compound (41) K salt having a benzotricarbocyanine structure and 6 sulfonic acid groups; compound (6) K salt, compound (4) K salt, compound (45) K salt, compound (31), compound (31) K salt, compound (3) K salt, compound (2) K salt, compound (43) K salt and compound (11) having a tricarbocyanine structure and 4-5 sulfonic acid groups; and control compounds ICG and NK-1967. Each test

compound was dissolved in distilled water (0.5 mg/ml) and used. The obtained each compound solution was administered from the tail vein of the mice (5.0 mg/kg). Blood was taken from the mice at 0.5, 1, 4 and 24 hours after the administration of the compounds and centrifuged to give plasma.

The fluorescence intensity of the plasma was measured by a spectrofluorescence meter (RF 5300 PC, SHIMADZU CORPORATION). A calibration curve of each compound was drawn and compound concentration in plasma was calculated. The results are shown in Fig. 10.

The compounds of the present invention remained in plasma at high concentration for a long time.

Experimental Example 5 : Acute toxicity

Reduction of toxicity by the introduction of sulfonic acid group and reduction thereof by conversion into sodium salt were studied.

The test compounds were those listed in Table 5.

Each test compound was dissolved in distilled water to give a compound solution. This solution was intravenously injected to the conscious mice from the tail vein. The mice were monitored for 3 days after the administration, and acute toxicity [LD_{50} (mg/kg body weight)] was estimated. The results are shown in Table 5.

Table 5

Number of sulfonic acid group	Compound	LD ₅₀ (mg/kg body weight)
Three or more	compound (11)K salt	350
	compound (11)	1980
	compound (31)K salt	350
	compound (31)	>3550
	compound (31)Ca salt	2000
	compound (31) pyridinium salt	1000-2000
	compound (45)K salt	550
	compound (45)	1100-1220
	compound (43)K salt	300-350
	compound (43)	1630
	compound (41)K salt	470
	compound (41)	>1010
	compound (29)K salt	470
	compound (29)	>1010
	compound (54)	>5000
	compound (6)K salt	350
	compound (3)K salt	530
	compound (4)K salt	450
	compound (2)K salt	610
Two or less	ICG	70
	NK1967	20


An increase in the number of sulfonic acid group in a molecule or conversion to a sodium salt resulted in striking reduction of acute toxicity.

The near infrared fluorescent contrast agent of the present invention is excited by an excitation light and emits near infrared fluorescence. This infrared fluorescence is superior in transmission through biological tissues. Thus, detection of lesions in the deep part of a living body has been made possible. In addition, the inventive contrast agent is superior in water solubility and low toxic, and therefore, it can be used safely.


PCT/EP99/07088

WHAT IS CLAIMED IS

1. A near infrared fluorescent contrast agent comprising a compound having three or more sulfonic acid groups in a molecule, which is represented by the formula [I]




wherein R^1 and R^2 are the same or different and each is a substituted or unsubstituted alkyl; Z^1 and Z^2 are each nonmetallic atoms necessary for forming a substituted or unsubstituted condensed benzo ring or condensed naphtho ring; r is 0, 1 or 2; $L^1 - L^7$ are the same or different and each is a substituted or unsubstituted methine, provided that when r is 2, L^6 and L^7 that occur in duplicate are the same or different; and X and Y are the same or different and each is a group of the formula







wherein R^3 and R^4 are the same or different and each is substituted or unsubstituted alkyl, or a pharmaceutically acceptable salt thereof.


2. The near infrared fluorescent contrast agent of claim 1, which is free of a carboxylic acid group in a molecule.
3. The near infrared fluorescent contrast agent of claim 1 or claim 2, wherein, in the formula [I], r is 1.
4. The near infrared fluorescent contrast agent of any of claims 1 to 3, wherein

- 4 or more sulfonic acid groups are contained in a molecule.
5. The near infrared fluorescent contrast agent of any of claims 1 to 4, wherein 10 or less sulfonic acid groups are contained in a molecule.
6. The near infrared fluorescent contrast agent of any of claims 1 to 4, wherein 8 or less sulfonic acid groups are contained in a molecule.
7. The near infrared fluorescent contrast agent of any of claims 1 to 6, wherein the pharmaceutically acceptable salt is a sodium salt.
8. The near infrared fluorescent contrast agent of any of claims 1 to 7, which is for tumor imaging.
9. The near infrared fluorescent contrast agent of any of claims 1 to 7, which is for angiography.
10. A sodium salt of a compound of the formula [II] having three or more sulfonic acid groups in a molecule




wherein R^1 , R^2 , $L^1 - L^7$, X and Y are as defined above; and R^5 to R^{16} are the same or different and each is a hydrogen atom, a sulfonic acid group, a carboxyl group, a hydroxyl group, an alkyl(sulfoalkyl)amino group, a bis(sulfoalkyl)amino group, a sulfoalkoxy group, a (sulfoalkyl)sulfonyl group or a (sulfoalkyl)aminosulfonyl group, exclusive of the groups of the following formulas






and




11. The sodium salt of claim 10, wherein, in the formula [II], R^1 and R^2 are each a lower alkyl having 1 to 5 carbon atoms substituted by a sulfonic acid group and X and Y are the same or different and each is




wherein R^{17} and R^{18} are unsubstituted lower alkyl having 1 to 5 carbon atoms.


12. The sodium salt of claim 11, having the formula








13. A sodium salt of a compound of the formula [III-1] having three or more sulfonic acid groups in a molecule




wherein L^1-L^7 are as defined above, R^{19} and R^{20} are lower alkyl having 1 to 5 carbon atoms and substituted by sulfonic acid group, $R^{21} - R^{28}$ are the same or different and each is hydrogen atom, sulfonic acid group, carboxyl group, hydroxyl group, alkyl(sulfoalkyl)amino group, bis(sulfoalkyl)amino group, sulfoalkoxy group, (sulfoalkyl)sulfonyl group or (sulfoalkyl)aminosulfonyl group, and X' and Y' are the same or different and each is a group of the formula



wherein R^{17} and R^{18} are as defined above, exclusive of the groups of the following formulas




and




14. The sodium salt of claim 13, wherein, in the formula [III-1], L₄ is methine substituted by alkyl having 1 to 4 carbon atoms.

15. The sodium salt of claim 13, which is a sodium salt of the compound of the formula [III-2] having three or more sulfonic acid groups in a molecule




wherein R¹⁹ – R²⁸, X' and Y' are as defined above, Z³ is a non-metallic atom group necessary for forming a 5- or 6-membered ring and A is hydrogen atom or a monovalent group.

16. The sodium salt of claim 15, having the formula



17. The sodium salt of claim 13, having the formula



18. The sodium salt of any of claims 10, 11, 13, 14 and 15, comprising 4 or more sulfonic acid groups in a molecule.

19. The sodium salt of any of claims 10, 11, 13, 14, 15 and 18, comprising 10 or less sulfonic acid groups in a molecule.

20. The sodium salt of any of claims 10, 11, 13, 14, 15 and 18, comprising 8 or less sulfonic acid groups in a molecule.





21. A near infrared fluorescent contrast agent comprising the sodium salt of any of claims 10 to 20.

22. The near infrared fluorescent contrast agent of claim 21, which is for tumor imaging.


23. The near infrared fluorescent contrast agent of claim 21, which is for angiography.

24. A method of fluorescence imaging comprising introducing the near infrared fluorescent contrast agent of claim 1 into a living body, exposing the body to an excitation light, and detecting near infrared fluorescence from the contrast agent.




25. The sodium salt of claim 10, which is at least one member selected from the group consisting of the compounds of the following formulas











[illegible]












26. The sodium salt of claim 13, which is at least one member selected from the group consisting of the compounds of the following formulas



































and









27. The near infrared fluorescent contrast agent of claim 1, comprising at least one compound selected from the group consisting of the compounds of the following formulas





















































and



28. The sodium salt of claim 15, wherein the monovalent group of A is substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted aralkyl, lower alkoxy, optionally substituted substituted amino, alkylcarbonyloxy, substituted or unsubstituted alkylthio, substituted or unsubstituted arylthio, cyano, nitro or halogen atom.




Fig. 2




Fig. 4




Fig. 1




Fig. 3

TOSTED 462260




Fig. 5

09/787394

Fig. 7




Fig. 9



Fig. 6



Fig. 8









Fig. 10

FIG. 10

FIG. 11











Fig. 14

T03T03"166.62.60

**DECLARATION FOR PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

NEAR INFRARED FLUORESCENT CONTRAST AGENT AND FLUORESCENCE IMAGING

the specification of which

☐ is attached hereto

☒ was filed on 16 SEPTEMBER 1999 as United States Application Number or PCT International Application Number PCT/EP99/07088 and (if applicable) was amended on _____

I hereby authorize our attorneys to insert the serial number assigned to this application.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR §1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate, or §365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 USC §119			
APPLICATION NO.	COUNTRY	DAY/MONTH/YEAR FILED	PRIORITY CLAIMED
10/283301	JAPAN	18/09/1998	YES

I hereby claim the benefit under 35 U.S.C. §119(e) of any United States provisional application(s) listed below.

PROVISIONAL APPLICATION(S) UNDER 35 U.S.C. §119(e)	
APPLICATION NUMBER	FILING DATE

I hereby claim the benefit under 35 U.S.C. §120 of any United States application, or §365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. §112.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR §1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

PRIOR U.S./PCT INTERNATIONAL APPLICATION(S) DESIGNATED FOR BENEFIT UNDER 37 U.S.C. §120		
APPLICATION NO.	FILING DATE	STATUS — PATENTED, PENDING, ABANDONED

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith: I. William Millen (19,544); John L. White (17,746); Anthony J. Zelano (27,969); Alan E.J. Branigan (20,565); John R. Moses (24,983); Harry B. Shubin (32,004); Brion P. Heaney (32,542); Richard J. Traverso (30,595); John A. Sopp (33,103); Richard M. Lebovitz (37,067); John H. Thomas (33,460); Nancy J. Axelrod (44,014); James E. Ruland (37,432) and Jennifer J. Branigan (40,921)

Correspondence Address:
MILLEN WHITE ZELANO & BRANIGAN, P.C.
Suite 1400
2200 Clarendon Boulevard
Arlington, VA 22201
TEL (703) 243-6333
FAX (703) 243-6410



PATENT & TRADEMARK OFFICE

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of sole or first inventor (given name, family name)	
Naoto MIWA	
Signature <i>Naoto Miwa</i>	Date <i>March 10, 2001</i>
Residence <i>JPX</i> Tokyo, Japan	Citizenship Japan
Post Office Address 1-21-2-2707, Miyamachi, Fuchu-shi, Tokyo 183-0023 Japan	
Full Name of additional joint inventor (given name, family name)	
Michihito INAGAKI	
Signature <i>Michihito Inagaki</i>	Date <i>April 24, 2001</i>
Residence <i>JPX</i> Osaka, Japan	Citizenship Japan
Post Office Address 16-10-303, Minoh 3-chome, Minoh-shi, Osaka 562-0001 Japan	
Full Name of additional joint inventor (given name, family name)	
Hiroaki EGUCHI	
Signature <i>Hiroaki Eguchi</i>	Date <i>April 23, 2001</i>
Residence <i>JPX</i> Hyogo, Japan	Citizenship Japan
Post Office Address 1-72, Seiwadainishi 2-chome, Kawanishi-shi, Hyogo 666-0143 Japan	
Full Name of additional joint inventor (given name, family name)	
Masafumi OKUMURA	
Signature <i>Masafumi Okumura</i>	Date <i>April 27, 2001</i>
Residence <i>JPX</i> Shiga, Japan	Citizenship Japan
Post Office Address 4-21, Chino 1-chome, Otsu-shi, Shiga 520-0111 Japan	
Full Name of additional joint inventor (given name, family name)	
Yoshio INAGAKI	
Signature <i>Yoshio Inagaki</i>	Date <i>March 15, 2001</i>
Residence <i>JPX</i> Kanagawa, Japan	Citizenship Japan
Post Office Address Fuji Photo Film Co., Ltd., 210, Naganuma, Minamiashigara-shi, Kanagawa 250-0193 Japan	
Full Name of additional joint inventor (given name, family name)	
Toru HARADA	
Signature <i>Toru Harada</i>	Date <i>March 15, 2001</i>
Residence <i>JPX</i> Kanagawa, Japan	Citizenship Japan
Post Office Address Fuji Photo Film Co., Ltd., 210, Naganuma, Minamiashigara-shi, Kanagawa 250-0193 Japan	

K:\PAT\Sch\1799\SCH 1799 DEC.wpd